

Klein Factors in multiple Fractional Quantum Hall edge tunneling

R. Guyon¹, P. Devillard^{1,2}, T. Martin^{1,3} and I. Safi¹

¹ Centre de Physique Théorique, Case 907 Luminy, 13288 Marseille Cedex 9, France

² Université de Provence, 13331 Marseille Cedex 03, France

³ Université de la Méditerranée, 13288 Marseille Cedex 9, France

A fractional quantum Hall liquid with multiple edges is considered. The computation of transport quantities such as current, noise and noise cross correlations in such multiple edge samples requires the implementation of so called Klein factors, which insure the correct quasiparticle exchange properties. The commutation relations of these factors are obtained by closing the system into a single edge. The non-equilibrium Green's function formalism associated with such factors is derived for a simple Laughlin fraction of the Hall effect. It is shown explicitly how Klein factors enter the calculation of the noise cross correlations, as well as the correction to the Poisson limit for the noise.

The edge state picture of the fractional quantum Hall effect [1] (FQHE) has led to fundamental discoveries. Various systems have been studied to analyze the quasiparticle excitations which tunnel through such quantum fluids [2,3]. In particular, for a point contact geometry with two edges, the detection of the backscattering noise leads to the measurement of the quasiparticle charge [4]. At the same time, resonant antidot tunneling geometries have allowed a capacitive measurement of the charge [5] in multiple edge geometries. Aside from this situation, multi edge geometries have not received much attention. Yet it was argued recently [6] that the detection of noise correlations in a three edge system constitutes a direct link to the quasiparticles statistics. One edge can be depleted due to the tunneling of quasiparticles into other edges. The Hanbury-Brown and Twiss type geometry [7] of Ref. [6] probes whether two quasi-holes on the injecting edge are allowed to overlap. Here, the key point is that quasiparticle edge tunneling operators need to be supplemented by Klein factors in order to specify the statistics. Klein factors were so far mostly introduced to tackle fermionic problems [8], yet it seems that their implementation is quite relevant for complex fractional Hall geometries.

The purpose here is to give a pedagogical view of how fractional statistics enters these multiple edge geometries: which (fractional) phase is generated when two quasiparticles from different edges are exchanged? We focus on Laughlin filling factors $\nu = (2p + 1)^{-1}$ (p integer). First, specifying a quasiparticle tunneling Hamiltonian, we derive the algebra which is obeyed by Klein factors. Next, we introduce the Keldysh Green's functions for the Klein factors. These tools are then applied to study the non-equilibrium current and noise correlations in a multi-edge system. In particular, it will be shown that Klein factors are unnecessary for simple, two edge samples. Finally, we argue that when considering the non-equilibrium average of a tunneling operator between two edges, there remains always a contribution associated with Klein factors, which is symptomatic of quasiparticle exchange with other edge “reservoirs”.




FIG. 1. Multiple edge geometry: tunneling processes T_i between the injecting edge specified by ϕ_0 and other edges, with fields labelled $\phi_1, \phi_2, \dots, \phi_N$. Curvilinear abscissas ($s, s'_1, s'_2, \dots$) are attached to each edge. In our disconnected edge geometry only the full lines defining the edges are kept. When considering the connected, single edge problem the dotted lines are added to fix the topological constraints. Inside frame shows how the transport properties in T_2 are affected by the fluctuating current in T_1 .

Our starting point is the Hamiltonian [9] which describes a system of decoupled edges $H_0 = (v_F \hbar / 4\pi) \sum_{l=0}^N \int ds (\partial_s \phi_l)^2$ with s the curvilinear abscissa. The fields $\partial_s \phi_l$ enter the description of long wave length excitations along each edge.

Tunneling between edge 0 and the other edges is described by the hopping Hamiltonian [3]:

$$H_t = \sum_{l=1}^N \sum_{\epsilon=\pm} \Gamma_l e^{-i\epsilon e^* \chi_l / \hbar c} T_l^{(\epsilon)} \quad (1)$$

where $T_l = T_l^{(+)} = \psi_0^\dagger(s_l) \psi_l(s'_l) = T_l^{(-)\dagger}$. $\psi_l^\dagger$ creates a quasiparticle excitation with charge $e^* = \nu e$ on edge l . Here “0” refers to a specific edge where, for instance, current can be injected into the others. A voltage bias can be imposed as usual using the Peierls substitution, i.e. $e^* \chi_l(t) = \hbar \omega_0 c t$ for a constant DC

bias $V_l = \hbar\omega_0/e^*$ imposed between edges 0 and l . The quasiparticle operators are written in a bosonized form $\psi_l(s) = (2\pi\alpha)^{-1/2}e^{-i\sqrt{\nu}\phi_l(s)}$ with α a short distance cutoff. Note that no fractional statistics is taken into account when quasiparticles from different edges are exchanged, whereas [10]:

$$\psi_l(s)\psi_l(s') = e^{\pm i\sqrt{\nu}\pi\text{sgn}(s-s')}\psi_l(s')\psi_l(s) \quad (2)$$

on a given edge. Fractional statistics is a consequence of the Laughlin correlated state describing the FQHE. The commutation relations for the ψ 's should originate from the derivation Chern–Simons boundary action. Here, we conjecture that the commutation relations are similar to those for a system of connected edges. For this reason, Klein factors are introduced, which means that each quasiparticle operator ψ_l is changed to $F_l\psi_l$. The requirement of unitarity, $F_l^\dagger F_l = F_l F_l^\dagger = 1$, is consistent with the fermionic case ($\nu = 1$).

Consider a closed system (Fig.1) where edges are connected with an open contour which brings all the edges into one. Klein factors are not necessary in this system because the commutation relations of tunneling operators are enforced by the single chiral bosonic field ϕ . Yet, by looking at the commutation relations of different tunneling operators we show below that for a system with the same topology but where edges are disconnected, the Klein factors can be computed. The operator describing tunneling from edge “0” (location s_l) to edge “ l ” (location s'_l) reads $T_l = (2\pi\alpha)^{-1}e^{i\sqrt{\nu}\phi(s_l)}e^{-i\sqrt{\nu}\phi(s'_l)}$.

Using the Baker–Campbell–Hausdorff formula [11], one can compute the commutator of two tunnel operators:

$$[T_j, T_k] = e^{-i\pi\nu}T_kT_j \left[1 - e^{-i\pi\nu[\text{sgn}(s_j-s_k)+\text{sgn}(s'_j-s'_k)]} \right] \quad (3)$$

Chirality imposes that $\text{sgn}(s_k - s_j) = -\text{sgn}(s'_j - s'_k)$ if the two tunneling paths do not cross. In this convention, which will be kept throughout the paper, $[T_j, T_k] = 0$. In the opposite situation where the tunneling paths have a crossing, $T_jT_k = e^{2i\pi\nu\text{sgn}(s_j-s_k)}T_kT_j$.

We now turn to the same system (same topology) depicted in Fig. 1 but where all edges are disconnected. In order to enforce fractional statistics, we introduce Klein factors in such a way that the tunneling operators become $T_l = \psi_0^\dagger(s_l)\psi_l(s'_l)F_0F_l^\dagger$. We require that the commutator of the tunneling operators of this disconnected system give the same results as the connected one. The Klein factors thus have to satisfy the relation :

$$(F_0F_k^\dagger)(F_0F_j^\dagger) = e^{i\nu\pi\text{sgn}(s_k-s_j)}(F_0F_j^\dagger)(F_0F_k^\dagger) \quad (4)$$

Eq. (4) is compatible with fractional statistics, i.e. $F_jF_k = e^{-i\pi\nu p_{jk}}F_kF_j$, for $j \neq k$ where $p_{jk} = \pm 1$. For arbitrary j and k , p_{jk} are the elements of an antisymmetric $N \times N$ matrix with elements 0, 1, -1 which are

tied to the edge configuration. Using Eq. (4), we find a connection between the elements p_{jk} and the sign of the algebraic distance : $\text{sgn}(s_k - s_j) = p_{k0} + p_{0j} + p_{jk}$. If for instance $k = 2$ and $j = 1$ with $\text{sgn}(s_2 - s_1) = 1$, we can choose : $p_{01} = -1 = -p_{02} = -p_{12}$.

These statistical constraints are now applied to a define problem of fractional Hall edge transport: the computation of current and noise correlations associated with several tunneling processes. We return to $N + 1$ edges in the previous system with N tunnelings paths. We represent the current on lead l using the Keldysh formulation of non-equilibrium transport:

$$\langle I_l \rangle = \frac{1}{2} \sum_{\eta=\pm} \langle T_K I_l(t^\eta) e^{-\frac{i}{\hbar} \int_K H_t(t') dt'} \rangle \quad (5)$$

where η labels the branch of the time ordering contour K .

The cross current fluctuations (i.e. the noise correlations) between two leads j and k are given by:

$$S_{jk}(t, t') = \sum_{\eta=\pm} \langle T_K I_j(t^\eta) I_k(t'^{-\eta}) e^{-\frac{i}{\hbar} \int_K H_t(t_1) dt_1} \rangle \quad (6)$$

The perturbative computation of these operators is achieved by expanding the evolution operator. The finite frequency noise $S_{jk}(\omega) = \sum_n S_{jk}^{(n)}(\omega)$ (n^{th} order in Γ_j 's) is the Fourier transform of the real time correlator (Eq. (6)). The $(n + 2)^{\text{th}}$ order of the noise is :

$$\begin{aligned} S_{jk}^{(n+2)}(\omega) = & \int dt e^{i\omega t} \left\{ \frac{\Gamma_j^{(\epsilon_j)} \Gamma_k^{(\epsilon_k)}}{n!} \left(-\frac{i}{\hbar} \right)^n \int dt_1 \dots dt_n \right. \\ & \times \sum_{\eta\eta_1 \dots \eta_n} \sum_{\epsilon\epsilon'} \sum_{\epsilon_1 \dots \epsilon_n} \epsilon\epsilon' e^{i\epsilon\omega_0 t - i\omega_0 \sum_{l=1}^n \epsilon_l t_l} \\ & \times \langle T_K T_j^{(\epsilon)}(t^\eta) T_k^{(\epsilon')} (t'^{-\eta}) \sum_{q_1=1}^N \Gamma_{q_1}^{(\epsilon_1)} T_{q_1}^{(\epsilon_1)}(t_1^{\eta_1}) \dots \\ & \dots \sum_{q_n=1}^N \Gamma_{q_n}^{(\epsilon_n)} T_{q_n}^{(\epsilon_n)}(t_n^{\eta_n}) \rangle - \langle I_j(t) \rangle \langle I_k(0) \rangle \left. \right\} \quad (7) \end{aligned}$$

The lowest non-vanishing contribution to the noise in a given terminal ($j = k$) is of order 2 in the tunneling amplitude, i.e. the Poisson result. The cross correlations require to compute the 4th order contribution. For example, the non-vanishing contribution to noise correlator between two terminals ($j < k$) :

$$\begin{aligned} S_{jk}^{(4)}(t, t') = & \frac{-2e^{*2}}{\hbar^2} \left(\frac{-i}{\hbar} \right)^2 \sum_{\eta\eta_1\eta_2} \sum_{\epsilon\epsilon'\epsilon_1\epsilon_2} \epsilon\epsilon' |\Gamma_j \Gamma_k|^2 \eta_1 \eta_2 \int dt_1 \int dt_2 \\ & \langle T_K T_j^{(\epsilon)}(t^\eta) T_k^{(\epsilon')} (t'^{-\eta}) T_j^{(\epsilon_1)}(t_1^{\eta_1}) T_k^{(\epsilon_2)}(t_2^{\eta_2}) \rangle \\ & \exp \left[i\epsilon\chi_j(t) + i\epsilon'\chi_k(t') + i \sum_{l=j,k} \epsilon_l \chi_l(t_l) \right] \quad (8) \end{aligned}$$

To this order, the Keldysh ordered product contains the product of two contributions. One of such originates from the computation of the average of the bosonic fields associated with each edge. This contains the dynamical aspect of these fields, and gives rise to exponentiated chiral Green functions. The other factor has no dynamics other than that specified by time ordering; It contains the Klein factors:

$$\begin{aligned} & \langle T_K \left\{ T_j^{(\epsilon)}(t^\eta) T_k^{(\epsilon')}(t'^{-\eta}) T_j^{(\epsilon_1)}(t_1^{\eta_1}) T_k^{(\epsilon_2)}(t_2^{\eta_2}) \right\} \rangle = \\ & (2\pi\alpha)^{-4} \langle T_K e^{-i\sqrt{\nu}\epsilon\phi_j^\eta(0,t)} e^{-i\sqrt{\nu}\epsilon_1\phi_j^{\eta_1}(0,t_1)} \rangle \\ & \times \langle T_K e^{-i\sqrt{\nu}\epsilon'\phi_k^{-\eta}(0,t')} e^{-i\sqrt{\nu}\epsilon_2\phi_k^{\eta_2}(0,t_2)} \rangle \\ & \times \langle T_K e^{+i\sqrt{\nu}\epsilon\phi_0^\eta(-s_{kj}/2,t)} e^{+i\sqrt{\nu}\epsilon'\phi_0^{-\eta}(s_{kj}/2,t')} \rangle \\ & \times \langle T_K e^{+i\sqrt{\nu}\epsilon_1\phi_0^{\eta_1}(-s_{kj}/2,t_1)} e^{+i\sqrt{\nu}\epsilon_2\phi_0^{\eta_2}(s_{kj}/2,t_2)} \rangle \\ & \times \langle T_K (F_0 F_j^\dagger)^{(\epsilon)}(t^\eta) (F_0 F_k^\dagger)^{(\epsilon')}(t'^{-\eta}) \rangle \\ & \times \langle T_K (F_0 F_j^\dagger)^{(\epsilon_1)}(t_1^{\eta_1}) (F_0 F_k^\dagger)^{(\epsilon_2)}(t_2^{\eta_2}) \rangle, \end{aligned} \quad (9)$$

where $s_{kj} = s_k - s_j$. The contraction of the chiral fields gives rise to quasiparticle conservation laws $\epsilon = -\epsilon_j$ and $\epsilon' = -\epsilon_k$. It is most convenient to compute the time ordered Klein factor product using a bosonization formulation similar to that used for the quasiparticle fields:

$$F_0 F_k^\dagger \equiv e^{-i\sqrt{\nu}\theta_k}. \quad (10)$$

Introducing the fact that $\text{sgn}(s_k - s_j) = 1$ in Eq. 4, we find (using the Hausdorff formula) that $[\theta_j, \theta_k] = i\pi$. From the fields θ_i , annihilation and creation operators a and $a^\dagger$ can then be defined as $a \equiv \frac{1}{\sqrt{2\pi}}(\theta_j + i\theta_k)$, with the usual property that $\langle a^\dagger a \rangle_0 = 0$ ("ground" state) and $\langle aa^\dagger \rangle = 1$.

The ground state expectation value of products of the bosonic fields reads: $\langle \theta_j \theta_k \rangle_0 = -\langle \theta_k \theta_j \rangle_0 = i\frac{\pi}{2}$ and $\langle \theta_j \theta_j \rangle = \langle \theta_k \theta_k \rangle = \frac{\pi}{2}$.

Because we are dealing with a non-equilibrium transport situation, a Keldysh matrix is introduced for the bosonic fields of Eq.(10) :

$$\begin{aligned} g_{jk}^{\eta_1\eta_2}(t) & \equiv \langle T_K \left\{ \theta_j(t^{\eta_1}) \theta_k(0^{\eta_2}) \right\} \rangle \\ & = i\frac{\pi}{2} \left\{ \frac{\eta_1 + \eta_2}{2} \text{sgn}(t) - \frac{\eta_1 - \eta_2}{2} \right\}. \end{aligned} \quad (11)$$

For the same tunneling events one finds $g_{ll}^{\eta_1\eta_2}(t) = \frac{\pi}{2}$.

The application of these Klein factor Green's function is most relevant when considering time ordered products

$$\begin{aligned} & \langle T_K \Pi_{l=j,k} e^{i\epsilon_{l1}\sqrt{\nu}\Theta_l(t_1^{\eta_{l1}})} \dots e^{i\epsilon_{lm_l}\sqrt{\nu}\Theta_l(t_{lm_l}^{\eta_{lm_l}})} \rangle \\ & = \exp\left(-\nu \sum_{l \leq l'} \sum_{k=1}^{m_l} \sum_{k'=1}^{m_{l'}} \epsilon_{lk} \epsilon_{l'k'} g_{ll'}^{\eta_{lk}\eta_{l'k'}}(t_{lk} - t_{l'k'})\right) \end{aligned} \quad (12)$$

$(l, l' = j, k)$ which vanishes unless $\sum_{k=1}^{m_l} \epsilon_{lk} = 0$.

Consider first the case of a two edge system, where only one Klein factor is necessary. In Eq. (12), the tunnel locations are $l = l' = 1$, so that the only Green's function

present is $g_{11}^{\eta_1\eta_2}(t) = \pi/2$. Together with the requirement of Eq. (12) to give a non-zero result, the term in the exponential then vanishes. This completes the proof that for a two edge system, when computing non-equilibrium averages for the current, the density fluctuations or the noise, the Klein factors do not contribute [12], and this, order by order in the perturbation series.

We now turn to the computation of the zero frequency current cross correlations between tunneling paths 1 and 2 in presence of a DC bias. Eq. (8) is integrated over t :

$$\begin{aligned} \tilde{S}_{jk}(\omega = 0) & = \frac{2e^{*2}}{\hbar^4} \sum_{\eta\eta_1\eta_2} \sum_{\epsilon\epsilon'} \epsilon\epsilon' \eta_1\eta_2 \int dt \int dt_1 \int dt_2 \\ & \frac{|\Gamma_j \Gamma_k|^2}{(2\pi\alpha)^4} e^{i\omega_0[\epsilon(t-t_1)-\epsilon't_2]} e^{2\nu[G_{\eta\eta_1}(0,t-t_1)+G_{-\eta\eta_2}(0,-t_2)]} \\ & \times \left(\frac{e^{\nu\epsilon\epsilon'[G_{\eta\eta_2}(-s_{kj},t-t_2)+G_{-\eta\eta_1}(s_{kj},-t_1)]}}{e^{\nu\epsilon\epsilon'[G_{\eta-\eta}(-s_{kj},t)+G_{\eta_1\eta_2}(-s_{kj},t_1-t_2)]}} e^{i\xi} - 1 \right) \end{aligned} \quad (13)$$

Where the Klein factor time ordered product reads :

$$e^{i\xi} = \langle T_K e^{-i\epsilon\sqrt{\nu}\theta_j(t^\eta)} e^{-i\epsilon'\sqrt{\nu}\theta_k(0^{-\eta})} e^{i\epsilon\sqrt{\nu}\theta_j(t_1^{\eta_1})} e^{i\epsilon'\sqrt{\nu}\theta_k(t_2^{\eta_2})} \rangle. \quad (14)$$

Note that the product of current averages $\langle I_j \rangle \langle I_k \rangle$ has to be subtracted before computing the Fourier transform. The resulting expression then identifies with the noise correlator which is typically measured in experiments [13]. Because of the symmetry properties of the Keldysh Greens function, only the configuration $\eta_1 = -\eta = -\eta_2$ needs to be considered and the configuration $\epsilon = -\epsilon'$ is chosen. Indeed, this turns out to be the relevant configuration which generates a series solution of the noise correlations [6]. Choosing the Klein factor correlator with $\epsilon = +$:

$$\begin{aligned} & \langle T_K e^{-i\sqrt{\nu}\theta_j(t^\eta)} e^{i\sqrt{\nu}\theta_k(0^{-\eta})} e^{i\sqrt{\nu}\theta_j(t_1^{\eta_1})} e^{-i\sqrt{\nu}\theta_k(t_2^{\eta_2})} \rangle = \\ & = \exp \left[\nu \left(g_{jk}^{\eta-\eta}(t) + g_{jj}^{\eta-\eta}(t-t_1) - g_{jk}^{\eta\eta}(t-t_2) \right) \right] \\ & \times \exp \left[\nu \left(-g_{kj}^{-\eta-\eta}(-t_1) + g_{kk}^{-\eta\eta}(-t_2) + g_{jk}^{-\eta\eta}(t_1-t_2) \right) \right] \\ & = \exp \left[\frac{i\nu\pi}{2} \eta (\text{sgn}(-t_1) - \text{sgn}(t-t_2)) \right] \end{aligned} \quad (15)$$

The Klein factor contribution with $\epsilon = -$ is simply the complex conjugate of Eq. (15). This term is crucial for a correct description of the noise cross correlations in the FQHE. For $\nu = 1$, it allows to recover the negative correlation result obtained with scattering theory [14]. For $\nu = 1/3$, the noise correlations exhibit an anti-bunching behavior, which is weaker than the integer case.

Finally, we turn to the computation of non-equilibrium averages attached to a given lead: the zero frequency current fluctuations in lead 1, denoted $\tilde{S}_j(\omega = 0)$. Perturbation theory tells us that there are two contributions to the current and noise:

$$\langle I_j \rangle = \langle I_j \rangle_0 + \langle I_j \rangle_K \quad (16)$$

$$\tilde{S}_j = \tilde{S}_j^0 + \tilde{S}_j^K \quad (17)$$

$\langle I_j \rangle_0$ and $\tilde{S}_j^0$ invoke only the tunneling operators T_j and $T_j^\dagger$ between edges 0 and j . As discussed above, these terms do not involve Klein factors as they couple only two edges, and they thus correspond exactly to the current and noise in a two terminal system derived in [15]:

$$\begin{aligned} \langle I_j \rangle_0 &\equiv -e\nu_F \int_{-\infty}^{+\infty} d\theta \rho(\theta) |S_{+-}(\theta - \theta_B)|^2 \\ \tilde{S}_j^0 &\equiv \frac{\nu e}{2(1-\nu)} \left[V \frac{d\langle I_j \rangle_0}{dV} - \langle I_j \rangle_0 \right] \end{aligned} \quad (18)$$

with $\rho(\theta)$ the kink density, $S_{+-}(\theta - \theta_B)$ the kink transmission probability and the barrier height appears via the kink parameter θ_B . Its leading order is proportional to $|\Gamma_j|^2$, the Poisson limit. On the other hand, the second contribution of Eq.(17) involves additional tunneling processes between the injecting edge 0 and other edges k , for instance $k > j$. In perturbation theory, these additional terms are proportional to $|\Gamma_j^p \Gamma_k^{p'}|^2$ ($p, p' \geq 1$) and they signal the possibility that quasiparticles from an external “reservoir” (edge k) are exchanged with the injecting edge 0. The leading order $|\Gamma_j \Gamma_k|^2$ correction to the Klein factor contribution reads:

$$\begin{aligned} \tilde{S}_j^K(\omega = 0) &= \frac{2|\Gamma_j \Gamma_k|^2}{h^4 \alpha^4} \sum_{\eta\eta_1\eta_2} \sum_{\epsilon\epsilon'} \eta_1 \eta_2 \int dt \int dt_1 \int dt_2 \\ &e^{i\omega_0[\epsilon t + \epsilon'(t_1 - t_2)]} e^{2\nu[G_{\eta-\eta_1}(0, t) + G_{\eta_1\eta_2}(0, t_1 - t_2)]} \\ &\times \left(\frac{e^{\nu\epsilon\epsilon'[G_{\eta\eta_2}(-a, t-t_2) + G_{-\eta\eta_1}(-a, -t_1)]}}{e^{\nu\epsilon\epsilon'[G_{\eta\eta_1}(-a, t-t_1) + G_{-\eta\eta_2}(-a, -t_2)]}} e^{i\xi'} - 1 \right) \end{aligned} \quad (19)$$

Where the Klein factor time ordered product reads:

$$\begin{aligned} e^{i\xi'} &= \langle T_K e^{-i\epsilon\sqrt{\nu}\theta_j(t^\eta)} e^{i\epsilon\sqrt{\nu}\theta_j(0^{-\eta})} e^{-i\epsilon'\sqrt{\nu}\theta_k(t_1^{\eta_1})} e^{i\epsilon'\sqrt{\nu}\theta_k(t_2^{\eta_2})} \rangle \\ &= e^{\nu\pi} e^{i\nu\pi\epsilon\epsilon'((\eta+\eta_2)\text{sgn}(t-t_2) + (-\eta+\eta_1)\text{sgn}(-t_1))/4} \\ &e^{i\nu\pi\epsilon\epsilon'(-(\eta+\eta_1)\text{sgn}(t-t_1) + (\eta-\eta_2)\text{sgn}(-t_2))/4} \end{aligned} \quad (20)$$

The calculation of the current and its contribution $\langle I_1 \rangle_K$ linked to the Klein factors follows the same guidelines, except that the Keldysh evolution operator in Eq. (5) is expanded to third order.

In conclusion, a systematic procedure is provided for computing transport quantities such as current, noise, noise correlations in multiple FQHE edges. Generalizations to hierarchical fractional Hall states and to multi-quasiparticle tunneling are envisioned. Multiple quantum Hall edge transport problems are necessary to address the nature of the statistics of the underlying excitations: the computation of the noise cross correlations requires Klein factors. Multiple edges allow to study fractional quantum transport through a constriction of a prepared fluctuating injecting current [16]. This injecting current fluctuates because of passage through a prior

gate, and the whole system requires once again three edges.

The current and the noise contain an “exact” contribution where all other tunneling processes are discarded, together with Klein factor corrections. This latter contribution illustrates how current and noise are affected by the presence of additional stochastic contributions. It is symptomatic of a tunneling process (T_2) which is influenced by another via a non linear coupling (T_1): in Fig. 1, the fluctuations at 1 propagate to the right along the bottom edge, affecting all transport properties at 2. In the language of quantum dissipative systems, it can be mapped to the problem of a brownian motion [17] in a periodic potential with two coupled particles.

T.M. acknowledges support from the LSF program at the Brown Submicron Center of Weizmann Institute.

-
- [1] R. B. Laughlin, Rev. Mod. Phys. **71**, 863 (1999); H.L. Stormer, *ibid.* 875 (1999).
 - [2] C. L. Kane and M. P. A. Fisher, Phys. Rev. Lett. **72**, 724 (1994).
 - [3] C. de C. Chamon, D. E. Freed, and X. G. Wen, Phys. Rev. B **51**, 2363 (1995-II).
 - [4] L. Saminadayar, D. C. Glatelli, Y. Jin, and B. Etienne, Phys. Rev. Lett. **79**, 2526 (1997); R. de-Picciotto *et al.*, Nature **389**, 162 (1997).
 - [5] V. J. Goldman and B. Su, Science **267**, 1010 (1995); V. J. Goldman, I. Karakurt, Jun Liu and A. Zaslavsky, cond-mat/0010093.
 - [6] I. Safi, P. Devillard and T. Martin, Phys. Rev. Lett. **86**, 4628 (2001).
 - [7] Hanbury-Brown and Q. R. Twiss, Nature **177**, 27 (1956).
 - [8] M. Fabrizio and A. Gogolin, Phys. Rev. B **51**, 17827 (1995); J. von Delft and H. Schoeller, Annalen der Physik **4**, 225 (1998); C. Nayak *et al.*, Phys. Rev. B **59**, 15694 (1999).
 - [9] X.G. Wen, Int. J. Mod. Phys. B **6**, 1711 (1992); Adv. Phys. **44**, 405 (1995).
 - [10] M.P.A. Fisher and L. I. Glazman in *Mesoscopic Electron Transport*, L. Kouwenhoven, G. Schön and L. Sohn eds., NATO ASI Series E345, Kluwer Ac. Publ., Dordrecht (1997).
 - [11] See, for instance, H. Kleinert, in *Path Integrals in Quantum Mechanics, Statistics and Polymer Physics*, pp. 139 (World Scientific 1995).
 - [12] S. Kettmann Phys. Rev. B. **55**, 2512 (1997).
 - [13] M. Henny *et al.*, Science **296**, 284 (1999).
 - [14] T. Martin and R. Landauer, Phys. Rev. B **45**, 1742 (1992); M. Büttiker, *ibid.* **46**, 12485 (1992).
 - [15] P. Fendley, A. Ludwig, and H. Saleur, Phys. Rev. Lett. **74**, 3005 (1995); *ibid.* **75**, 2196 (1995).
 - [16] M. Heiblum, private communication.
 - [17] A. Schmid, Phys. Rev. Lett. **51**, 1506 (1983); F. Guinea, V. Hakim and A. Muramatsu, *ibid.* **54**, 263 (1985).